



Swaziland

Multiple Indicator Cluster Survey 2014

Key Findings

May, 2015



TABLE OF CONTENTS

SWAZILAND MICS AT A GLANCE.....	1
CHILD MORTALITY.....	2
NUTRITION.....	3
CHILD HEALTH.....	6
WATER AND SANITATION.....	8
REPRODUCTIVE HEALTH.....	9
CHILD DEVELOPMENT	11
LITERACY AND EDUCATION.....	12
CHILD PROTECTION	13
HIV/AIDS AND SEXUAL BEHAVIOUR	15
ACCESS TO MASS MEDIA AND ICT.....	18
SUBJECTIVE WELL-BEING	18
TOBACCO AND ALCOHOL USE.....	19
NOTES	20

The Swaziland Multiple Indicator Cluster Survey (MICS5) was carried out in 2014 by the Central Statistical Office in collaboration with United Nations Children’s Fund (UNICEF), United Nations Population Fund (UNFPA) and United Nations, Educational, Scientific and Cultural Organization (UNESCO), as part of the global MICS programme. Technical support was provided by the United Nations Children’s Fund (UNICEF). The government of Swaziland, UNFPA, UNESCO and UNICEF provided financial support.

The global MICS programme was developed by UNICEF in the 1990s as an international household survey programme to collect internationally comparable data on a wide range of indicators on the situation of children and women. MICS surveys measure key indicators that allow countries to generate data for use in policies and programmes, and to monitor progress towards the Millennium Development Goals (MDGs) and other internationally agreed upon commitments.

The objective of this report is to facilitate the timely dissemination and use of results from the Swaziland Multiple Indicator Cluster Survey (MICS) prior to the release of full tables and the final survey report that will contain detailed information on all survey findings by various demographic, social, economic and cultural characteristics. The final survey report is expected to be released in the fourth quarter of 2015.

Results presented in this report are not expected to change and are considered final. For more information on indicators and the analysis conducted in the full final report please go to mics.unicef.org.

Suggested citation:

Central Statistical Office. 2015 Swaziland *Multiple Indicator Cluster Survey 2014, Key Findings*. Mbabane, Swaziland: Central Statistical Office

SWAZILAND MICS AT A GLANCE

Survey implementation			
Sample frame	Swaziland Population and Housing Census, 2007	Questionnaires	Household Women (age 15-49) Men (age 15-59) Children under five
Interviewer training	June – July 2014	Fieldwork	July – October 2014
Survey sample			
Households		Children under five	
- Sampled	5,211	- Eligible	2,728
- Occupied	4,981	- Mothers/caretakers interviewed	2,693
- Interviewed	4,865	- Response rate (Per cent)	98.7
- Response rate (Per cent)	97.7		
Women		Men	
- Eligible for interviews	5,001	- Eligible for interviews	1,629
- Interviewed	4,762	- Interviewed	1,459
- Response rate (Per cent)	95.2	- Response rate (Per cent)	89.6

Survey population			
Average household size	4.0	Percentage of population living in	
Percentage of population under:		- Urban areas	26.7
- Age 5	12.7	- Rural areas	73.3
- Age 18	46.6	- Hhohho	25.0
Percentage of women age 15-49 years with at least one live birth in the last 2 years	20.1	- Manzini	37.1
		- Shiselweni	17.9
		- Lubombo	20.0

Housing characteristics	
Percentage of households with	
- Electricity	65.0
- Finished floor	95.3
- Finished roofing	94.2
- Finished walls	88.8
Mean number of persons per room used for sleeping	2.10

Household or personal assets	
Percentage of households that own	
- A television	54.9
- A refrigerator	55.3
- Agricultural land	65.4
- Farm animals/livestock	56.8
Percentage of households where at least a member has or owns a	
- Mobile phone	95.9
- Car or truck	22.7


CHILD MORTALITY

Early childhood mortality

MICS Indicator	Indicator	Description	Value ^A
1.1	Neonatal mortality rate	Probability of dying within the first month of life	20
1.2	MDG 4.2 Infant mortality rate	Probability of dying between birth and the first birthday	50
1.3	Post-neonatal mortality rate	Difference between infant and neonatal mortality rates	30
1.4	Child mortality rate	Probability of dying between the first and the fifth birthdays	18
1.5	MDG 4.1 Under-five mortality rate	Probability of dying between birth and the fifth birthday	67

^A Indicator values are per 1,000 live births and refer to the five-year period before the survey

Figure 1: Early childhood mortality rates, Swaziland Multiple Indicator Cluster Survey, 2014




Note: Indicator values are per 1,000 live births

NUTRITION

Nutritional status ⁱ				
MICS Indicator	Indicator	Description	Value	
2.1a	MDG 1.8 Underweight prevalence (a) Moderate and severe (b) Severe	Percentage of children under age 5 who fall below		
2.1b		(a) minus two standard deviations (moderate and severe) (b) minus three standard deviations (severe) of the median weight for age of the WHO standard	5.8 1.6	
2.2a	Stunting prevalence (a) Moderate and severe (b) Severe	Percentage of children under age 5 who fall below		
2.2b		(a) minus two standard deviations (moderate and severe) (b) minus three standard deviations (severe) of the median height for age of the WHO standard	25.5 7.2	
2.3a	Wasting prevalence (a) Moderate and severe (b) Severe	Percentage of children under age 5 who fall below		
2.3b		(a) minus two standard deviations (moderate and severe) (b) minus three standard deviations (severe) of the median weight for height of the WHO standard	2.0 0.4	
2.4	Overweight prevalence	Percentage of children under age 5 who are above two standard deviations of the median weight for height of the WHO standard	9.0	

Figure 2: Underweight, stunted, wasted and overweight children under age 5 (moderate and severe), Swaziland Multiple Indicator Cluster Survey, 2014



Breastfeeding and infant feeding			
MICS Indicator	Indicator	Description	Value
2.5	Children ever breastfed	Percentage of women with a live birth in the last 2 years who breastfed their last live-born child at any time	92.1
2.6	Early initiation of breastfeeding	Percentage of women with a live birth in the last 2 years who put their last newborn to the breast within one hour of birth	48.3
2.7	Exclusive breastfeeding under 6 months	Percentage of infants under 6 months of age who are exclusively breastfed ⁱⁱ	63.8
2.8	Predominant breastfeeding under 6 months	Percentage of infants under 6 months of age who received breast milk as the predominant source of nourishment ⁱⁱⁱ during the previous day	70.2
2.9	Continued breastfeeding at 1 year	Percentage of children age 12-15 months who received breast milk during the previous day	47.8
2.10	Continued breastfeeding at 2 years	Percentage of children age 20-23 months who received breast milk during the previous day	7.6
2.11	Median duration of breastfeeding	The age in months when 50 percent of children age 0-35 months did not receive breast milk during the previous day	13.0
2.12	Age-appropriate breastfeeding	Percentage of children age 0-23 months appropriately fed ^{iv} during the previous day	45.3
2.13	Introduction of solid, semi-solid or soft foods	Percentage of infants age 6-8 months who received solid, semi-solid or soft foods during the previous day	89.5
2.14	Milk feeding frequency for non-breastfed children	Percentage of non-breastfed children age 6-23 months who received at least 2 milk feedings during the previous day	52.8
2.15	Minimum meal frequency	Percentage of children age 6-23 months who received solid, semi-solid and soft foods (plus milk feeds for non-breastfed children) the minimum number of times ^v or more during the previous day	81.2
2.16	Minimum dietary diversity	Percentage of children age 6-23 months who received foods from 4 or more food groups ^{vi} during the previous day	62.4
2.17a	Minimum acceptable diet	(a) Percentage of breastfed children age 6-23 months who had at least the minimum dietary diversity and the minimum meal frequency during the previous day	48.6
2.17b		(b) Percentage of non-breastfed children age 6-23 months who received at least 2 milk feedings and had at least the minimum dietary diversity not including milk feeds and the minimum meal frequency during the previous day	29.7
2.18	Bottle feeding	Percentage of children age 0-23 months who were fed with a bottle during the previous day	31.7

Salt iodization

MICS Indicator	Indicator	Description	Value
2.19	Iodized salt consumption	Percentage of households with salt testing 15 parts per million or more of iodate	66.1


Low-birthweight

MICS Indicator	Indicator	Description	Value
2.20	Low-birthweight infants	Percentage of most recent live births in the last 2 years weighing below 2,500 grams at birth	8.0
2.21	Infants weighed at birth	Percentage of most recent live births in the last 2 years who were weighed at birth	92.3

CHILD HEALTH

Vaccinations			
MICS Indicator	Indicator	Description	Value
3.1	Tuberculosis immunization coverage	Percentage of children age 12-23 months who received BCG vaccine by their first birthday	97.5
3.2	Polio immunization coverage	Percentage of children age 12-23 months who received the third dose of OPV vaccine (OPV3) by their first birthday	83.9
3.3	Diphtheria, pertussis and tetanus (DPT3), (HepB3) and (Hib3) immunization coverage	Percentage of children age 12-23 months who received the third dose of DPT vaccine (DPT3) by their first birthday	90.1
3.4 MDG 4.3	Measles immunization coverage	Percentage of children age 12-23 months who received measles vaccine by their first birthday	89.3
3.8	Full immunization coverage	Percentage of children age 12-23 months who received all ^{vii} vaccinations recommended in the national immunization schedule by their first birthday	70.7

Figure 3: Vaccinations by age 12 months (measles by 24 months), Swaziland Multiple Indicator Cluster Survey, 2014



Tetanus toxoid

MICS Indicator	Indicator	Description	Value
3.9	Neonatal tetanus protection	Percentage of women age 15-49 years with a live birth in the last 2 years who were given at least two doses of tetanus toxoid vaccine within the appropriate interval prior to the most recent birth	83.0

Diarrhoea

MICS Indicator	Indicator	Description	Value
-	Children with diarrhoea	Percentage of children under age 5 with diarrhoea in the last 2 weeks	16.4
3.10	Care-seeking for diarrhoea	Percentage of children under age 5 with diarrhoea in the last 2 weeks for whom advice or treatment was sought from a health facility or provider	71.2
3.11	Diarrhoea treatment with oral rehydration salts (ORS) and zinc	Percentage of children under age 5 with diarrhoea in the last 2 weeks who received ORS and zinc	42.3
3.12	Diarrhoea treatment with oral rehydration therapy (ORT) and continued feeding	Percentage of children under age 5 with diarrhoea in the last 2 weeks who received ORT (ORS packet, pre-packaged ORS fluid, recommended homemade fluid or increased fluids) and continued feeding during the episode of diarrhoea	46.4

Acute Respiratory Infection (ARI) symptoms

MICS Indicator	Indicator	Description	Value
-	Children with ARI symptoms	Percentage of children under age 5 with ARI symptoms in the last 2 weeks	9.8
3.13	Care-seeking for children with ARI symptoms	Percentage of children under age 5 with ARI symptoms in the last 2 weeks for whom advice or treatment was sought from a health facility or provider	59.6
3.14	Antibiotic treatment for children with ARI symptoms	Percentage of children under age 5 with ARI symptoms in the last 2 weeks who received antibiotics	26.6

Solid fuel use


MICS Indicator	Indicator	Description	Value
3.15	Use of solid fuels for cooking	Percentage of household members in households that use solid fuels as the primary source of domestic energy to cook	62.3

WATER AND SANITATION

Water and sanitation				
MICS Indicator	Indicator	Description		Value
4.1	MDG 7.8	Use of improved drinking water sources	Percentage of household members using improved sources of drinking water	72.0
4.2		Water treatment	Percentage of household members in households using unimproved drinking water who use an appropriate treatment method	16.8
4.3	MDG 7.9	Use of improved sanitation	Percentage of household members using improved sanitation facilities which are not shared	53.0
4.4		Safe disposal of child's faeces	Percentage of children age 0-2 years whose last stools were disposed of safely	55.3

The data on place for handwashing and availability of soap is still being analyzed and will be presented in the final report

Figure 4: Use of improved water and sanitation in urban and rural areas, Swaziland Multiple Indicator Cluster Survey, 2014



REPRODUCTIVE HEALTH


Contraception and unmet need

MICS Indicator	Indicator	Description	Value
-	Total fertility rate	Total fertility rate ^A for women age 15-49 years	3.3
5.1	MDG 5.4 Adolescent birth rate	Age-specific fertility rate ^A for women age 15-19 years	87
5.2	Early childbearing	Percentage of women age 20-24 years who had at least one live birth before age 18	16.7
5.3	MDG 5.3 Contraceptive prevalence rate	Percentage of women age 15-49 years currently married or in union who are using (or whose partner is using) a (modern or traditional) contraceptive method	66.1
5.4	MDG 5.6 Unmet need	Percentage of women age 15-49 years who are currently married or in union who are fecund and want to space their births or limit the number of children they have and who are not currently using contraception	15.2

^AThe age-specific fertility rate is defined as the number of live births to women in a specific age group during a specified period, divided by the average number of women in that age group during the same period, expressed per 1,000 women. The age-specific fertility rate for women age 15-19 years is also termed as the adolescent birth rate.

The total fertility rate (TFR) is calculated by summing the age-specific fertility rates calculated for each of the 5-year age groups of women, from age 15 through to age 49. The TFR denotes the average number of children to which a woman will have given birth by the end of her reproductive years (by age 50) if current fertility rates prevailed.

Figure 5: Use of contraception, Swaziland Multiple Indicator Cluster Survey, 2014



Modern methods		65.5
Female sterilization		3.7
Male sterilization		0.0
IUD		0.7
Injectables		21.9
Implants		3.8
Pill		10.5
Male condom		23.9
Female condom		0.8
Diaphragm/Foam/Jelly		0.2

Traditional methods		0.6
Periodic abstinence		0.2
Withdrawal		0.3
Other		0.1

Any method	66.1
-------------------	-------------

No method	33.9
------------------	-------------


Maternal and newborn health

MICS Indicator	Indicator	Description	Value
5.5a	MDG 5.5	Antenatal care coverage Percentage of women age 15-49 years with a live birth in the last 2 years who were attended during their last pregnancy that led to a live birth (a) at least once by skilled health personnel (b) at least four times by any provider	98.5
5.5b	MDG 5.5		76.1
5.6	Content of antenatal care	Percentage of women age 15-49 years with a live birth in the last 2 years who had their blood pressure measured and gave urine and blood samples during the last pregnancy that led to a live birth	90.0
5.7	MDG 5.2	Skilled attendant at delivery Percentage of women age 15-49 years with a live birth in the last 2 years who were attended by skilled health personnel during their most recent live birth	88.3
5.8	Institutional deliveries	Percentage of women age 15-49 years with a live birth in the last 2 years whose most recent live birth was delivered in a health facility	87.7
5.9	Caesarean section	Percentage of women age 15-49 years whose most recent live birth in the last 2 years was delivered by caesarean section	11.6

Post-natal health checks

MICS Indicator	Indicator	Description	Value
5.10	Post-partum stay in health facility	Percentage of women age 15-49 years who stayed in the health facility for 12 hours or more after the delivery of their most recent live birth in the last 2 years	90.2
5.11	Post-natal health check for the newborn	Percentage of last live births in the last 2 years who received a health check while in facility or at home following delivery, or a post-natal care visit within 2 days after delivery	90.1
5.12	Post-natal health check for the mother	Percentage of women age 15-49 years who received a health check while in facility or at home following delivery, or a post-natal care visit within 2 days after delivery of their most recent live birth in the last 2 years	87.2

Figure 6: Place of delivery and post-natal health checks, Swaziland Multiple Indicator Cluster Survey, 2014




CHILD DEVELOPMENT

Child development			
MICS Indicator	Indicator	Description	Value
6.1	Attendance to early childhood education	Percentage of children age 36-59 months who are attending an early childhood education programme	29.5
6.2	Support for learning	Percentage of children age 36-59 months with whom an adult has engaged in four or more activities to promote learning and school readiness in the last 3 days	38.6
6.3	Father's support for learning	Percentage of children age 36-59 months whose biological father has engaged in four or more activities to promote learning and school readiness in the last 3 days	1.8
6.4	Mother's support for learning	Percentage of children age 36-59 months whose biological mother has engaged in four or more activities to promote learning and school readiness in the last 3 days	16.4
6.5	Availability of children's books	Percentage of children under age 5 who have three or more children's books	5.9
6.6	Availability of playthings	Percentage of children under age 5 who play with two or more types of playthings	67.0
6.7	Inadequate care	Percentage of children under age 5 left alone or in the care of another child younger than 10 years of age for more than one hour at least once in the last week	16.5
6.8	Early child development index	Percentage of children age 36-59 months who are developmentally on track in at least three of the following four domains: literacy-numeracy, physical, social-emotional, and learning	64.9

LITERACY AND EDUCATION

Literacy and education ^{viii}				
MICS Indicator	Indicator	Description	Value	
7.1	MDG 2.3	Literacy rate among young people	Percentage of young people age 15-24 years who are able to read a short simple statement about everyday life or who attended secondary or higher education (a) women (b) men	95.3 92.1
7.2		School readiness	Percentage of children in first grade of primary school who attended pre-school during the previous school year	60.6
7.3		Net intake rate in primary education	Percentage of children of school-entry age who enter the first grade of primary school	89.3
7.4	MDG 2.1	Primary school net attendance ratio (adjusted)	Percentage of children of primary school age currently attending primary or secondary school	97.7
7.5		Secondary school net attendance ratio (adjusted)	Percentage of children of secondary school age currently attending secondary school or higher	50.4
7.6	MDG 2.2	Children reaching last grade of primary	Percentage of children entering the first grade of primary school who eventually reach last grade	92.9
7.7		Primary completion rate	Number of children attending the last grade of primary school (excluding repeaters) divided by number of children of primary school completion age (age appropriate to final grade of primary school)	90.9
7.8		Transition rate to secondary school	Number of children attending the last grade of primary school during the previous school year who are in the first grade of secondary school during the current school year divided by number of children attending the last grade of primary school during the previous school year	85.5
7.9	MDG 3.1	Gender parity index (primary school)	Primary school net attendance ratio (adjusted) for girls divided by primary school net attendance ratio (adjusted) for boys	1.01
7.10	MDG 3.1	Gender parity index (secondary school)	Secondary school net attendance ratio (adjusted) for girls divided by secondary school net attendance ratio (adjusted) for boys	1.19

Figure 7: Education indicators by sex, Swaziland Multiple Indicator Cluster Survey, 2014



Note: All indicator values are in per cent

CHILD PROTECTION


Birth registration

MICS Indicator	Indicator	Description	Value
8.1	Birth registration	Percentage of children under age 5 whose births are reported registered	53.5

Child discipline

MICS Indicator	Indicator	Description	Value
8.3	Violent discipline	Percentage of children age 1-14 years who experienced psychological aggression or physical punishment during the last one month	88.3

Figure 8: Child disciplining methods, children age 1-14 years, Swaziland Multiple Indicator Cluster Survey, 2014



Early marriage and polygyny

MICS Indicator	Indicator	Description	Value
8.4	Marriage before age 15	Percentage of people who were first married or in union before age 15 (a) Women 15 – 49 years (b) Men 15 – 49 years (c) Men 15 – 59 years	1.3 0.2 0.2
8.5	Marriage before age 18	Percentage of people who were first married or in union before age 18 (a) Women 20 – 49 years (b) Men 20 – 49 years (c) Men 20 – 59 years	8.8 1.7 2.0
8.6	Young people age 15-19 years currently married or in union	Percentage of young people age 15-19 years who are married or in union (a) Women (b) Men	4.0 0.0
8.7	Polygyny	Percentage of people who are in a polygynous union (a) Women 15 – 49 years (b) Men 15 – 49 years (c) Men 15 – 59 years	11.7 8.2 10.4
8.8a 8.8b	Spousal age difference	Percentage of young women who are married or in union and whose spouse is 10 or more years older, (a) among women age 15-19 years, (b) among women age 20-24 years	(32.5) 22.5

() Figures that are based on 25-49 unweighted cases

Attitudes towards domestic violence

MICS Indicator	Indicator	Description	Value
8.12	Attitudes towards domestic violence	Percentage of people who state that a husband is justified in hitting or beating his wife in at least one of the following circumstances: (1) she goes out without telling him, (2) she neglects the children, (3) she argues with him, (4) she refuses sex with him, (5) she burns the food (a) Women 15 – 49 years (b) Men 15 – 49 years (c) Men 15 – 59 years	19.9 17.0 16.3

Children's living arrangements

MICS Indicator	Indicator	Description	Value
8.13	Children's living arrangements	Percentage of children age 0-17 years living with neither biological parent	33.2
8.14	Prevalence of children with one or both parents dead	Percentage of children age 0-17 years with one or both biological parents dead	20.4
8.15	Children with at least one parent living abroad	Percentage of children 0-17 years with at least one biological parent living abroad	13.0

HIV/AIDS AND SEXUAL BEHAVIOUR

HIV/AIDS knowledge and attitudes			
MICS Indicator	Indicator	Description	Value
-	Have heard of AIDS	Percentage of people who have heard of AIDS (a) Women 15 – 49 years (b) Men 15 – 49 years (c) Men 15 – 59 years	99.8 99.8 99.8
9.1	MDG 6.3 Knowledge about HIV prevention among young people	Percentage of young people age 15-24 years who correctly identify ways of preventing the sexual transmission of HIV ^{ix} , and who reject major misconceptions about HIV transmission ^x (a) Women (b) Men	49.1 50.9
9.2	Knowledge of mother-to-child transmission of HIV	Percentage of people who correctly identify all three means ^{xi} of mother-to-child transmission of HIV (a) Women 15 – 49 years (b) Men 15 – 49 years (c) Men 15 – 59 years	66.5 52.9 52.2
9.3	Accepting attitudes towards people living with HIV	Percentage of people expressing accepting attitudes on all four questions ^{xii} toward people living with HIV (a) Women 15 – 49 years (b) Men 15 – 49 years (c) Men 15 – 59 years	37.4 36.2 37.0
9.S1	Discriminatory attitudes Towards people living with HIV	Percentage of people who report discriminatory attitudes towards people living with HIV ^{xiii} (a) Women 15 – 49 years (b) Men 15 – 49 years (c) Men 15 – 59 years	37.4 36.2 37.0

HIV testing			
MICS Indicator	Indicator	Description	Value
9.4	People who know where to be tested for HIV	Percentage of people who state knowledge of a place to be tested for HIV (a) Women 15 – 49 years (b) Men 15 – 49 years (c) Men 15 – 59 years	97.4 96.4 96.5
9.5	People who have been tested for HIV and know the results	Percentage of people who have been tested for HIV in the last 12 months and who know their results (a) Women 15 – 49 years (b) Men 15 – 49 years (c) Men 15 – 59 years	66.5 54.5 55.3
9.6	Sexually active young people who have been tested for HIV and know the results	Percentage of young people age 15-24 years who have had sex in the last 12 months, who have been tested for HIV in the last 12 months and who know their results (a) Women (b) Men	80.2 62.3
9.7	HIV counselling during antenatal care	Percentage of women age 15-49 years who had a live birth in the last 2 years and received antenatal care during the pregnancy of their most recent birth, reporting that they received counselling on HIV during antenatal care	89.9
9.8	HIV testing during antenatal care	Percentage of women age 15-49 years who had a live birth in the last 2 years and received antenatal care during the pregnancy of their most recent birth, reporting that they were offered and accepted an HIV test during antenatal care and received their results	95.3

Sexual behaviour

MICS Indicator	Indicator	Description	Value
9.9	Young people who have never had sex	Percentage of never married young people age 15-24 years who have never had sex (a) Women (b) Men	54.5 53.6
9.10	Sex before age 15 among young people	Percentage of young people age 15-24 years who had sexual intercourse before age 15 (a) Women (b) Men	3.0 2.8
9.11	Age-mixing among sexual partners	Percentage of women age 15-24 years who had sex in the last 12 months with a partner who was 10 or more years older	14.5
9.12	Multiple sexual partnerships	Percentage of people who had sexual intercourse with more than one partner in the last 12 months (a) Women 15 – 49 years (b) Men 15 – 49 years (c) Men 15 – 59 years	3.3 21.1 20.6
9.13	Condom use at last sex among people with multiple sexual partnerships	Percentage of people who report having had more than one sexual partner in the last 12 months who also reported that a condom was used the last time they had sex (a) Women 15 – 49 years (b) Men 15 – 49 years (c) Men 15 – 59 years	66.0 82.6 80.8
9.14	Sex with non-regular partners	Percentage of sexually active young people age 15-24 years who had sex with a non-marital, non-cohabitating partner in the last 12 months (a) Women (b) Men	37.3 40.9
9.15	MDG 6.2 Condom use with non-regular partners	Percentage of young people age 15-24 years reporting the use of a condom during the last sexual intercourse with a non-marital, non-cohabiting sex partner in the last 12 months (a) Women (b) Men	70.9 93.4

Orphans

MICS Indicator	Indicator	Description	Value
9.16	MDG 6.4 Ratio of school attendance of orphans to school attendance of non-orphans	Proportion attending school among children age 10-14 years who have lost both parents divided by proportion attending school among children age 10-14 years whose parents are alive and who are living with one or both parents	1.00

Male circumcision

MICS Indicator	Indicator	Description	Value
9.17	Male circumcision	Percentage of men who reported have been circumcised (a) Men age 15 – 49 years (b) Men age 15 – 59 years	25.0 24.3

ACCESS TO MASS MEDIA AND ICT

Access to mass media

MICS Indicator	Indicator	Description	Value
10.1	Exposure to mass media	Percentage of people who at least once a week, read a newspaper or magazine, listen to the radio, and watch television	
		(a) Women 15 - 49 years	22.6
		(b) Men 15 - 49 years	30.9
		(c) Men 15 - 59 years	30.1

Use of information/communication technology

MICS Indicator	Indicator	Description	Value
10.2	Use of computers	Percentage of young people age 15-24 years who used a computer during the last 12 months	
		(a) Women	42.7
		(b) Men	48.9
10.3	Use of internet	Percentage of young people age 15-24 years who used the internet during the last 12 months	
		(a) Women	39.8
		(b) Men	55.6

SUBJECTIVE WELL-BEING

Subjective well-being

MICS Indicator	Indicator	Description	Value
11.1	Life satisfaction	Percentage of young people age 15-24 years who are very or somewhat satisfied with their life, overall	
		(a) Women	82.5
		(b) Men	84.8
11.2	Happiness	Percentage of young people age 15-24 years who are very or somewhat happy	
		(a) Women	75.5
		(b) Men	75.6
11.3	Perception of a better life	Percentage of young people age 15-24 years whose life improved during the last one year, and who expect that their life will be better after one year	
		(a) Women	62.2
		(b) Men	62.8

TOBACCO AND ALCOHOL USE

Tobacco use

MICS Indicator	Indicator	Description	Value
12.1	Tobacco use	Percentage of people who smoked cigarettes, or used smoked or smokeless tobacco products at any time during the last one month	
		(a) Women 15 – 49 years	1.3
		(b) Men 15 – 49 years	14.7
		(c) Men 15 – 59 years	15.1
12.2	Smoking before age 15	Percentage of people who smoked a whole cigarette before age 15	
		(a) Women 15 – 49 years	0.2
		(b) Men 15 – 49 years	2.9
		(c) Men 15 – 49 years	2.8

Alcohol use

MICS Indicator	Indicator	Description	Value
12.3	Use of alcohol	Percentage of people who had at least one alcoholic drink at any time during the last one month	
		(a) Women 15 – 49 years	5.5
		(b) Men 15 – 49 years	25.5
		(c) Men 15 – 59 years	25.9
12.4	Use of alcohol before age 15	Percentage of people who had at least one alcoholic drink before age 15	
		(a) Women 15 – 49 years	1.5
		(b) Men 15 – 49 years	5.9
		(c) Men 15 – 59 years	5.8

NOTES

ⁱWeight and height/length measurements were successfully completed for 98.7 and 98.4 per cent of children under age 5, respectively

ⁱⁱInfants receiving breast milk, and not receiving any other fluids or foods, with the exception of oral rehydration solution, vitamins, mineral supplements and medicines

ⁱⁱⁱInfants receiving breast milk and certain fluids (water and water-based drinks, fruit juice, ritual fluids, oral rehydration solution, drops, vitamins, minerals, and medicines), but do not receive anything else (in particular, non-human milk and food-based fluids)

^{iv}Infants age 0-5 months who are exclusively breastfed, and children age 6-23 months who are breastfed and ate solid, semi-solid or soft foods

^vBreastfeeding children: Solid, semi-solid, or soft foods, two times for infants age 6-8 months, and three times for children 9-23 months; Non-breastfeeding children: Solid, semi-solid, or soft foods, or milk feeds, four times for children age 6-23 months

^{vi}The indicator is based on consumption of any amount of food from at least 4 out of the 7 following food groups: 1) grains, roots and tubers, 2) legumes and nuts, 3) dairy products (milk, yogurt, cheese), 4) flesh foods (meat, fish, poultry and liver/organ meats), 5) eggs, 6) vitamin-A rich fruits and vegetables, and 7) other fruits and vegetables

^{vii} Full vaccination includes the following: BCG, Polio1, Polio2, Polio3, DPT1/HEPB1/ HIB1, DPT2/HEPB2/ HIB2, DPT3/HEPB3/ HIB3, Measles 1. NB: PCV was not included in the calculation because it was introduced before the commencement of the MICS data collection exercise.

^{viii} Education indicators, wherever applicable, are based on information on reported school attendance (at any time during the school year), as a proxy for enrolment.

^{ix}Using condoms and limiting sex to one faithful, uninfected partner

^xThe two most common misconceptions about HIV transmission are included in the indicator calculation: i) mosquito bites and ii) sharing food with someone with HIV

^{xi}Transmission during pregnancy, during delivery, and by breastfeeding

^{xii}People (1) who think that a female teacher who is HIV-positive and is not sick should be allowed to continue teaching, (2) who would buy fresh vegetables from a shopkeeper or vendor who is HIV-positive, (3) who would not want to keep secret that a family member is HIV-positive, and (4) who would be willing to care for a family member with AIDS in own home

^{xiii}People who (1) would not buy vegetables from a shopkeeper or vendor who is HIV positive and (2) think children living with HIV should not be able to attend school with children who are HIV negative